

 **Europäisches Patentamt**
European Patent Office
Office européen des brevets

(11) Publication number:

0 162 764
A1

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 85400832.3

(51) Int. Cl.⁴: **A 61 K 47/00**
A 61 K 31/22, A 61 K 31/23

(22) Date of filing: 26.04.85

(30) Priority: 02.05.84 US 606054

(43) Date of publication of application:
27.11.85 Bulletin 85/48

(84) Designated Contracting States:
CH DE FR GB IT LI NL

(71) Applicant: **MERCK & CO. INC.**
126, East Lincoln Avenue P.O. Box 2000
Rahway New Jersey 07065(US)

(72) Inventor: **Alexander, Jose**
2909 Westdale
Lawrence Kansas 66044(US)

(72) Inventor: **Fix, Joseph A.**
824, Mississippi
Lawrence Kansas 66044(US)

(74) Representative: **Ahner, Francis et al,**
CABINET REGIMBEAU 26, avenue Kléber
F-75116 Paris(FR)

(54) Composition and method for enhancing absorption of drugs from gastrointestinal tract using acylcarnitines.

(57) Acylcarnitines used as drug absorption enhancing
agents for orally and rectally administered drugs.

EP 0 162 764 A1

- 1 -

IX-115

TITLE OF THE INVENTION

COMPOSITION AND METHOD FOR ENHANCING ABSORPTION OF
DRUGS FROM GASTROINTESTINAL TRACT USING ACYLCARNITINES.

5 BACKGROUND OF THE INVENTION

The invention relates to a novel method and
compositions for enhancing absorption of drugs from
the gastrointestinal tract by incorporating therein
an acylcarnitine absorption enhancing agent.

10

DESCRIPTION OF THE PRIOR ART

Although numerous references describe
carnitine-acylcarnitine translocase system in
mitochondria, the instant use of acylcarnitines to
15 promote gastrointestinal drug absorption constitutes
a novel and unobvious use of said acylcarnitine
compounds. The carnitine-acylcarnitine translocase

20

system has been localized in skeletal muscle, cardiac muscle, epididymal tissue, liver, kidney and brain. Evidence for the existence of this system in the non-muscle mucosal cells of the gastrointestinal tract, which is the barrier to drug absorption, has not been found. The system is located on the inner membrane of the mitochondria, an intracellular organelle and its functions are to transport fatty acids across the mitochondria membrane and lead to intraorganelle concentration and oxidation of the fatty acids. There is no evidence to indicate that acylcarnitines will promote transmembrane movement of another target molecule. Carnitine can transverse membranes, however, carnitine alone has no effect in the instant application. The acylcarnitine ester is required for the absorption promoting effect. Thus, the intramitochondrial transport of fatty acids, which is a known function of carnitine, is unrelated to its function as a gastrointestinal drug absorption promoter.

The use of acylcarnitines to promote gastrointestinal absorption affords several advantages over the prior art's non-related absorption promoting compounds. The acylcarnitines, especially these with medium chain fatty acid components, are more potent than the presently used absorption promoting agents. As an example, in aqueous solutions, the acylcarnitines are effective absorption promoting agents at levels as low as 0.05%. By contrast, the effective dose of other known absorption promoters is significantly higher: sodium salicylate - 1%, surfactants - 1%, chelating agents - 2%. This difference in potency affords opportunities for

reducing the required size of the dosage form and potentially minimizing side effects. The acylcarnitines cause reversible changes in gastrointestinal permeability to the target drug, indicating that a permanent change has not occurred. Other promoting agents, such as the surfactants, cause a relatively permanent change in gastrointestinal permeability, which is only overcome by turnover of the mucosal cells, a process which may require days for completion. By contrast, removal of acylcarnitines from the gastrointestinal tract results in reversion to normal permeability properties in less than 2 hours. This provides a significant advantage in that a rapid and reversible increase in drug absorption does not allow prolonged intervals during which potentially toxic or otherwise harmful agents might also be absorbed. Another potential advantage of the acylcarnitines is that, unlike chelating agent such as EDTA, the acylcarnitines may not necessarily sequester divalent cations (Mg^{++} or Ca^{++}) which are necessary for the normal functioning of cells. In other words, there is no tissue damage at concentrations of acylcarnitines which significantly increase drug absorption. In contrast to this, studies have indicated that surfactant activity, as with sodium lauryl sulfate, is generally associated with some degree of cellular damage. This lack of tissue damage affords a significant advantage to the use of acylcarnitines in promoting gastrointestinal drug absorption which can be metabolized through normal pathways in the body. This eliminates a potential problem from introducing substances which

are not normally present in the biochemical pathways of the body (e.g. salicylates, EDTA etc.)

SUMMARY OF THE INVENTION

5 It has been found that when poorly absorbed drugs are administered orally or rectally, the bioavailability of said drugs is increased by administering together with an acylcarnitine absorption enhancing agent.

10 Accordingly, it is an object of this invention to enhance the bioavailability of poorly absorbed drugs administered orally or rectally by administering therewith an acylcarnitine absorption enhancing agent.

15 It is an object of the invention to provide a new dosage form utilizing a novel class of acylcarnitines which when administered orally or rectally will provide an increased blood level of the therapeutic agent.

20 Another object of the invention is to provide an acylcarnitine absorption promoter of gastrointestinal drug absorption at concentrations which do not alter the normal morphology of the mucosal cells.

25 Still another object of the invention is to provide an acylcarnitine series of absorption agents that are endogenous and can be metabolized through normal pathways available in the body.

30 Other objects, features and advantages of the invention will be apparent to those skilled in the art from the detailed description of the invention which follows.


All of the foregoing objects are readily attained by providing a method and drug form wherein oral and rectal absorption of poorly absorbed drugs is enhanced. The method comprises the steps of
5 preparing a dosage form suitable for oral or rectal delivery, and a dosage form comprising an effective unit dosage amount of the poorly absorbed drug, an acylcarnitine absorption agent or pharmaceutically acceptable salt thereof, the agent being present in
10 said dosage form in an amount sufficient to be effective in enhancing the rate of the oral and rectal absorption of the therapeutic agent, and pharmaceutically acceptable excipients.

15 DETAILED DESCRIPTION OF THE INVENTION

The present invention, generally comprises the steps of preparing a dosage form capable of being orally or rectally administered, wherein the dosage form comprises an effective unit dosage amount of a
20 poorly absorbed drug and an acylcarnitine absorption enhancing agent, the acylcarnitine agent being present in the dosage form in a sufficient quantity to be effective in enhancing oral and rectal absorption rates and administering the dosage form to
25 warm-blooded animals. The amount of poorly absorbed drug varies over a wide range, but generally the therapeutically effective unit dosage amount of the selected poorly absorbed drug depends on that amount known in the art to obtain the desired results.

30 The compounds that are used as absorption enhancers in our method and drug forms are acylcarnitines which are β -acyloxy derivatives of

γ -trimethylaminobutyric acid and pharmaceutically acceptable salts thereof of the formula:



wherein R is:

- a) $\text{C}_2\text{-C}_{20}$ saturated acyl such as acetyl,
10 hexanoyl, octanoyl, lauroyl, myristoyl, palmitoyl, stearoyl and the like;
 - b) $\text{C}_2\text{-C}_{20}$ acyl with 1 to 6 double bonds such
as 2-hexenoyl, 9-decenoyl, 9-hexadecenoyl
(palmitoleoyl), oleoyl, myristoleoyl,
15 9,12-hexadecadienoyl, α -linoleoyl, γ -linolenoyl, arachidyl and the like;
 - c) $\text{C}_2\text{-C}_{20}$ hydroxyacyl with 1 to 3 hydroxy
groups such as 2-hydroxylauroyl, 2-hydroxymyristoyl,
2-hydroxypalmitoyl and the like;
 - 20 d) $\text{C}_4\text{-C}_{20}$ ketoacyl such as 6-ketodecanoyl, 4-keto-9,11,13-octadecatrienoyl and the like;
 - e) $\text{C}_5\text{-C}_{20}$ unsaturated hydroxyacyl such as
2-hydroxy-12-octadecenoyl and the like;
 - 25 f) $\text{C}_5\text{-C}_{20}$ carbalkoxyacyl such as ω -ethoxy-carbonyloctanoyl and the like;
- and X is a pharmaceutically acceptable counterion
such as chloride, sulfate, nitrate, perchlorate,
bromide, phosphate, acetate, benzoate, tartrate,
citrate, propionate, gluconate, lactate, maleate,
30 fumarate, bezylate, camsylate, esylate, gluceptate, mesylate, napsylate and the like.

The preferred oral and rectal absorption enhancing agents of the above formula are:

1. Acetylcarnitine
2. Hexanoylcarnitine
- 5 3. Lauroylcarnitine
4. Octanoylcarnitine
5. Myristoylcarnitine
6. Palmitoylcarnitine
7. Stearoylcarnitine
- 10 8. 2-Hexenoylcarnitine
9. 9-Decenoylcarnitine
10. 9-Hexadecenoylcarnitine
11. α -Linoleoylcarnitine
12. 2-Hydroxylauroylcarnitine
- 15 13. 2-Hydroxymyristoylcarnitine
14. 6-Keto-decanoylcarnitine
15. 12-Hydroxy-12-octadecenoylcarnitine
16. ω -Ethoxylcarbonyloctanoylcarnitine
17. 2-Hydroxypalmitoylcarnitine.

20 The most preferred absorption enhancing agents useful in our method and dosage forms are:

1. Acetylcarnitine
2. Hexanoylcarnitine
- 25 3. Octanoylcarnitine
4. Lauroylcarnitine
5. Myristoylcarnitine
6. Palmitoylcarnitine
7. Stearoylcarnitine.

30

The acylcarnitine absorption enhancing agents employed in the practice of this invention are

known compounds which are commercially available and processes for their preparation are disclosed throughout the art.

Various active agents provide beneficial
5 effects when administered to patients. Such agents which can be made more useful by enhancing its absorption in accordance with this invention, are exemplified by, but not limited to, the following classes of drugs:

10 1) β -lactam antibiotics such as cefoxitin, N-formamidinylthienamycin, amoxicillin, ampicillin, azlocillin, bacampicillin, carbenicillin, cefaclor, cefadroxil, cefamandole, cefaparole, cefatrizine, cefazoline, cefonicid, cefaperazone, ceforanide,
15 cefotaxime, cefotiam, cefroxadine, cefsulodin, ceftazidime, ceftizoxime, cephalaxin, cephaloglycin, cephaloridine, cephradine, cyclacillin, cloxacillin, dicloxacillin, floxacillin, hetacillin, methicillin, nafcillin, oxacillin, sarmoxacillin, sarcipillin,
20 talampicillin, ticaricillin, penicillin G, penicillin V, pivampicillin, piperacillin, pirbenicillin and the like.

2) Aminoglycoside antibiotics such as gentamycin, amikacin, astromicin, betamicin, butikacin,
25 butirosin, clindamycin, josamycin, kanamycin, neomycin, netilmicin, tobramycin and the like.

3) Antiviral agents such as ara C (cytarabine), acyclovir, floxuridine, ribavirin, vidarabine, idoxuridine, trifluridine and the like.

30 4) Amino acids such as methyl dopa, carbidopa, levodopa, fludalamine, α -aminobutyric acid and the like.

5) Smooth muscle relaxants such as theophylline, aminophylline, diphylline, oxtriphylline, ambu-
phylline, fenethylline, guathylline, pentoxifylline,
xanthinol niacinate, theophylline glycinate,
5 glucophylline and the like.

6) Polypeptides such as cyclo(N-Ala-Tyr-D-Trp-Lys-Val-Phe)acetate (363,586), somatostatin, insulin, gastrin, caerulein, cholecystokinin and the like.

10 7) Anti-inflammatory agents such as indomethacin, sulindac, ibuprofen and the like.

8) Diuretics such as aldactone, hydrochlorothiazide, amiloride, amiloride and hydrochloride and the like.

15 The enhancement of drug absorption in accordance with this invention is not by any means limited to the above drugs, but are in general applicable to other classes of drugs such as analgesics, anabolics, androgens, anorexics,
20 adrenergics, antiadrenergics, antiallergics, antibacterials, anticholinergics, antidepressants, antidiabetics, antifungal agents, antihypertensives, antineoplastics, antipsychotics, sedatives, cardiovascular agents, antiulcer agents, anti-
25 coagulants, anthelmintics, radio-opaques, radio-nuclide diagnostic agents and the like.

Generally, the amount of adjuvant employed in the practice of the invention ranges from 0.05-500 mg in each unit dose. The percentage of adjuvant in
30 the total combination of drug plus adjuvant is 0.05-50% with a preferred ratio of adjuvant in the total combination of adjuvant plus drug being 0.05-25%.

For oral administration, the formulations may be prepared as liquids, suspensions, capsules, tablets, coated tablets, and other standard procedures known in the art. The preferred formulation is a
5 compressed tablet composed of a minimum of 1 mg acylcarnitine ester with the pharmacologically required dose of drug and sufficient excipients to formulate an acceptable composition. For rectal application, the formulations may be prepared as
10 microenemas, suppositories, rectal tablets, and other standard procedures known in the art. The preferred formulation is a solid suppository composed of a minimum of 1 mg acylcarnitine ester with the pharmacologically required dose of drug and sufficient
15 suppository base to formulate an acceptable composition. The methods and choice of excipients and suppository bases are well known to those skilled in the art and the composition of said formulations is not limited to compressed tablets or solid suppositories by this invention.
20

The following examples illustrate preparation of various compositions of the invention. The examples should be construed as illustrations rather than limitations thereof.

25

EXAMPLE 1

Effect of palmitoylcarnitine chloride on the rectal absorption of drug entities. Experiments were performed with rats wherein each animal received an
30 aqueous microenema applied to the rectal cavity. The microenemas contained target drug entity (amount shown in table) in the presence or absence of 5 mg

0162764

- 11 -

IX-115

palmitoylcarnitine. Blood levels were monitored and the amount of drug absorbed calculated against intravenous administration and expressed as percent bioavailability.

5

10

15

20

25

30

Target Drug	Dose	Drug Class	Percent Bioavailability (mean \pm SD)	
			Control	with Palmitoylcarnitine-Cl
Sodium cefoxitin	2.5 mg	β -lactam antibiotic	2 \pm 1.1	68 \pm 7.2
Gentamicin sulfate	2.5 mg	aminoglycoside antibiotic	4 \pm 1.2	42 \pm 5.2
Cytarabine	2.5 mg	anti-viral anti-neoplastic	0.5 \pm 0.1	9 \pm 0.6
Theophylline	2.5 mg	smooth muscle relaxant	75 \pm 3.3	75 \pm 2.6
**#-363,586	0.1 mg	polypeptide	7 \pm 3.5	100 \pm 28.2
Methyldopa	2.5 mg	cardiovascular anti-hypertensive	6 \pm 0.3	100 \pm 8.9

** cyclo(N-Me-Ala-Tyr-D-Trp-Lys-Val-Phe)acetate

EXAMPLE 2

Concentration dependence of palmitoyl-
carnitine effect on rectal absorption of sodium
cefoxitin, a β -lactam antibiotic. Each animal
 5 received an aqueous microenema at pH 6 containing 2.5
 mg sodium cefoxitin in the presence or absence of
 varying amounts of palmitoylcarnitine. Blood samples
 were collected and assayed for sodium cefoxitin. The
 amount of drug absorbed is expressed as percent
 10 bioavailability as compared to intravenous
 administration.

15	Dose of Palmitoylcarnitine-Cl (mg)	Sodium Cefoxitin Percent Bioavailability mean \pm SD
	0	2 \pm 1.1
	0.025	1 \pm 0.4
	0.125	22 \pm 3.6
20	0.25	43 \pm 8.0
	1.25	81 \pm 13.1
	2.5	67 \pm 9.0
	5.0	68 \pm 7.2

25

EXAMPLE 3

Effect of various carnitine esters on rectal
absorption of sodium cefoxitin, a β -lactam antibiotic,
and cyclo(N-Ala-Tyr-D-Trp-Lys-Val-Phe)acetate, a
 30 polypeptide. Each animal received an aqueous
 microenema, pH 6, containing 2.5 mg sodium cefoxitin
 or 0.1 mg cyclo(N-Ala-Tyr-D-Trp-Lys-Val-Phe)-

acetate and 5.0 mg various carnitine esters of the general formula of this invention. Blood samples were collected and sodium cefoxitin or cyclo(N-Ala-Tyr-D-Trp-Lys-Val-Phe)acetate assayed.

5 The amount of drug absorbed is expressed as percent bioavailability versus intravenous administration.

		Percent Bioavailability (mean \pm SD)	
<u>Carnitine Ester</u>		<u>Sodium Cefoxitin*</u>	<u>**</u>
10	None	2 \pm 1.1	7 \pm 3.5
	Acetylcarnitine-Cl	6 \pm 3.4	39 \pm 21.9
	Hexanoylcarnitine-Cl	2 \pm 1.3	33 \pm 5.6
	Octanoylcarnitine-Cl	7 \pm 2.8	3 \pm 1.1
	Lauroylcarnitine-Cl	51 \pm 8.7	23 \pm 2.5
15	Myristoylcarnitine-Cl	27 \pm 2.1	100 \pm 45.0
	Palmitoylcarnitine-Cl	68 \pm 7.2	100 \pm 28.2
	Stearoylcarnitine-Cl	52 \pm 2.7	26 \pm 5.1

* β -lactam antibiotic

20 ** Cyclo(N-Ala-Tyr-D-Trp-Lys-Val-Phe)acetate

EXAMPLE 4

Importance of carnitine-palmitoyl ester linkage in absorption promoting effects on rectal sodium cefoxitin absorption. Each animal received an aqueous microenema, pH 6, containing 2.5 mg sodium cefoxitin in the presence or absence of 5.0 mg palmitic acid, carnitine or palmitoylcarnitine. Blood was collected and sodium cefoxitin measured.

25

30 Absorption of sodium cefoxitin is expressed as percent bioavailability versus intravenous administration.

		Sodium Cefoxitin
		Percent Bioavailability
		(mean \pm SD)
Compound		
5	None	2 \pm 1.1
	Palmitic acid	1 \pm 0.8
	Carnitine	1 \pm 0.6
	Palmitoylcarnitine-Cl	68 \pm 7.2

10

EXAMPLE 5

Reversibility of absorption promoting effect of carnitine esters/histological evaluation of mucosal surface. Two separate experiments demonstrate that carnitine esters cause no permanent change in rectal mucosal tissue at concentrations which effectively increase drug absorption. In experiment A, animals are treated with 5.0 mg palmitoylcarnitine alone and then tested for sodium cefoxitin absorption either immediately or after 1 hour or 2 hours of recovery. Sodium cefoxitin absorption is expressed as percent bioavailability and indicates the reversibility of the absorption promoting effect upon removal of the carnitine ester. In experiment B, animals are exposed to varying concentrations of palmitoylcarnitine for 20 minutes, and the tissue examined histologically for changes in responses to treatment.

30

	<u>A</u>		<u>B</u>	
	Interval Between Administra- tions (min)	Sodium Cef- oxitin Per- cent Bio- availability (mean \pm SD)	Palmitoyl- carnitine Dose (mg)	Histo- logical Evaluation
5				
10	0	68 \pm 7.2	5.0	Negligible morpho- logical changes
	60	36 \pm 9.2	2.5	normal
	120	9 \pm 6.6	1.25	normal
			0.25	normal
			0.125	normal
15			0.025	normal

EXAMPLE 6

Effect of palmitoylcarnitine chloride on
small intestinal absorption of polypeptides.
 Experiments were performed with rats wherein each
 animal received an aqueous solution applied to the
 duodenal region. The solutions contained target drug
 entity 0.1 mg. [cyclo(N-Ala-Tyr-D-Trp-Lys-Val-Phe)
 acetate] in the presence or absence of 5 mg
 palmitoylcarnitine. Blood levels were monitored and
 the amount of polypeptide absorbed calculated against
 intravenous administration and expressed as percent
 bioavailability.

0162764

- 17 -

IX-115

		Percent Bioavailability
Adjuvant		(mean \pm SD)
None		1
5	Palmitoylcarnitine-Cl	11 \pm 2.7

10

15

20


25

30

WHAT IS CLAIMED IS:

1. A pharmaceutical composition for enhancing gastrointestinal tract absorption of an orally or rectally administered formulation comprising a therapeutically effective dosage amount of a poorly absorbed drug and an acylcarnitine absorption enhancing agent of the formula:

10



- wherein R is saturated acyl(C₂-C₂₀), acyl-(C₂-C₂₀) with 1 to 6 double bonds, hydroxyacyl-(C₂-C₂₀) with 1 to 3 hydroxy groups, ketoacyl-(C₄-C₂₀), unsaturated hydroxyacyl(C₅-C₂₀) or carbalkoxyacyl(C₅-C₂₀) and x is a pharmaceutically acceptable counterion.

20

2. The composition of Claim 1 wherein the drug is selected from the group consisting of β-lactam antibiotics, aminoglycosides, antiviral agents, amino acids, smooth muscle relaxants, polypeptides, anti-inflammatory agents and diuretics and acylcarnitine absorption enhancing agent is selected from the group consisting of acetyl-carnitine, hexanoylcarnitine, lauroylcarnitine, octanoylcarnitine, myristoylcarnitine, palmitoylcarnitine, stearoylcarnitine, 2-hexenoylcarnitine, 9-decenoylcarnitine, 9-hexadecenoylcarnitine, α-linoleoylcarnitine, 2-hydroxylauroylcarnitine, 2-hydroxymyristoylcarnitine, 6-keto-decanoyl-

30


carnitine, 12-hydroxy-12-octadecenoylcarnitine, ω -ethoxycarbonyloctanoylcarnitine and 2-hydroxy-palmitoylcarnitine.

5 3. The composition of Claim 2 wherein the
drug is an Antibiotic selected from the group
consisting of cefoxitin, ampicillin, cefamanodole,
cefazoline, cefotaxime, cefaclor, cefalothin,
10 cefitizoxime, amoxicillin, N-formamidinylthienamycin,
cefadroxil, gentamycin, neomycin, clindamycin,
astromicin and betamycin; an Antiviral agent selected
from the group consisting of cytarabine, acylclovir,
trifluridine and vidarabine; an Amino acid selected
15 from the group consisting of methyldopa, carbidopa,
 α -aminobutyric acid and levadopa; a smooth Muscle
relaxant selected from the group consisting of
theophylline, aminophylline, xanthinol niacinate,
glucophylline; a Polypeptide selected from the group
20 consisting of gastrin, somatostatin, insulin and
cyclo(N-Me-Ala-Tyr-D-Trp-Lys-Val-Phe)acetate;
an anti-inflammatory agent selected from the group
consisting of indomethacin and sulindac; and a
Diuretic selected from the group consisting of
25 amiloride, hydrochlorothiazide and aldactone and the
agent is selected from the group consisting of
acetylcarnitine, hexanoylcarnitine, octanoyl-
carnitine, lauroylcarnitine, myristoylcarnitine,
palmitoylcarnitine and stearoylcarnitine.

30 4. The composition of Claim 3 wherein the
drug is cefoxitin, gentamycin, cytarabine, methyldopa,
theophylline, cyclo(N-Ala-Tyr-D-Trp-Lys-Val-Phe)-
acetate, indomethacin or amiloride and the agent is
palmitoylcarnitine.

5. The composition of Claim 1 further comprising pharmaceutically acceptable excipients.

6. A method of enhancing the rate of gastrointestinal absorption of an orally or rectally administered drug, said method comprising the steps of preparing a dosage form capable of being orally or rectally absorbed, said dosage form comprising a therapeutically effective dosage amount of said drug and an acylcarnitine absorption enhancing agent of the formula:



15

wherein R is saturated acyl(C₂-C₂₀), acyl-(C₂-C₂₀) with 1 to 6 double bonds, hydroxyacyl-(C₂-C₂₀) with 1 to 3 hydroxy groups, ketoacyl-(C₄-C₂₀), unsaturated hydroxyacyl(C₅-C₂₀) or carbalkoxyacyl(C₅-C₂₀) and X is a pharmaceutically acceptable counterion.

7. The method of Claim 6 wherein the drug is selected from the group consisting of β-lactam antibiotics, aminoglycoside antibiotics, antiviral agents, Amino acids, Smooth muscle relaxants, Polypeptides, Anti-inflammatory agents and diuretics and the Acylcarnitine absorption enhancing agent is selected from the group consisting of acetyl-carnitine, hexanoylcarnitine, lauroylcarnitine, octanoylcarnitine, myristoylcarnitine, palmitoylcarnitine, stearoylcarnitine, 2-hexenoylcarnitine,

1734M/0967A

- 21 -

IX115

9-decenoylcarnitine, 9-hexadecenoylcarnitine,
α-linoleoylcarnitine, 2-hydroxylauoylcarnitine,
2-hydroxymyristoylcarnitine, 6-keto-decanoyl-
carnitine, 12-hydroxy-12-octadecenoylcarnitine,
5 methoxycarbonyloctanoylcarnitine, and 2-hydroxy-
palmitoylcarnitine.

8. The method of Claim 7 wherein the agent
is selected from the group consisting of acetyl-
10 carnitine, hexanoylcarnitine, octanoylcarnitine,
lauroylcarnitine, myristoylcarnitine,
palmitoylcarnitine and stearoylcarnitine.

15

20

25

30



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT
which under Rule 45 of the European Patent Convention
shall be considered, for the purposes of subsequent
proceedings, as the European search report

0162764

Application number

EP 85 40 0832

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
X	FR-A- 2 482 588 (SIGMA-TAU IND. FARMACEUTICHE RIUNITE S.P.A.) * Page 15, line 23 - page 16, line 7; claims 3-4 * ---	1-5	A 61 K 47/00 A 61 K 31/22 A 61 K 31/23
X	FR-A- 2 244 487 (KAWASHIMA, Y.) * Page 9, lines 1-20; claims 1-6 * ---	1-5	
A	FR-A- 2 388 556 (SIGMA-TAU IND. FARMACEUTICHE RIUNITE S.P.A.) * Page 14, lines 1-13; claim 1 * -----	1-5	
			TECHNICAL FIELDS SEARCHED (Int. Cl.4)
			A 61 K C 07 C
INCOMPLETE SEARCH			
<p>The Search Division considers that the present European patent application does not comply with the provisions of the European Patent Convention to such an extent that it is not possible to carry out a meaningful search into the state of the art on the basis of some of the claims.</p> <p>Claims searched completely: 1-5 Claims searched incompletely: Claims not searched: 6-8 Reason for the limitation of the search: Method for treatment of the human or animal body by surgery or therapy (See art. 52(4) of the European Patent Convention)</p>			
Place of search The Hague		Date of completion of the search 06-08-1985	Examiner BRINKMANN
CATEGORY OF CITED DOCUMENTS		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER: _____**

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.